



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore y_1 = z + x \sin z + \frac{x^2}{1.2} \frac{d}{dz} (\sin^2 z) + \frac{x^3}{1.2.3} \left(\frac{d}{dz} \right)^2 (\sin^3 z) + \text{etc.}$$

$$\begin{aligned} \therefore y_1 &= z + x \sin z + \frac{x^2}{1.2} \frac{d}{dz} \left(\frac{1 - \cos 2z}{2} \right) + \frac{x^3}{1.2.3} \left(\frac{d}{dz} \right)^2 \left(\frac{3 \sin z - \sin 3z}{4} \right) \\ &\quad + \frac{x^4}{1.2.3.4} \left(\frac{d}{dz} \right)^3 \left(\frac{3 - 4 \cos 2z + \cos 4z}{8} \right) + \text{etc.} \\ &= z + x \sin z + \frac{1}{2} x^2 \sin 2z + \frac{1}{8} x^3 (3 \sin 3z - \sin z). \end{aligned}$$

$$\begin{aligned} \therefore v &= m + e \sin m + \frac{1}{2} e^2 \sin 2m + \frac{1}{8} e^3 (3 \sin 3m - \sin m) \\ &\quad + \frac{1}{8} e^4 (2 \sin 4m - \sin 2m) + \text{etc.} = m + e \sin v. \end{aligned}$$

$$\therefore \sin v = \sin m + \frac{1}{2} e \sin 2m + \frac{1}{8} e^2 (3 \sin 3m - \sin m) + \frac{1}{8} e^3 (2 \sin 4m - \sin 2m) + \text{etc.}$$

To develop $(1 - e \cos v)$ in terms of m : Let $f(y_1) = 1 - e \cos y_1$, $f'(y_1) = e \sin y_1$.

$$\therefore 1 - e \cos y_1 = (1 - e \cos z) + x \sin z (e \sin z) + \frac{x^2}{1.2} \frac{d}{dz} (\sin^2 z \cdot e \sin z) + \text{etc.}$$

Performing the operations as before we get after substituting v for y_1 , m for z , e for x ,

$$\begin{aligned} 1 - e \cos v &= 1 - e \cos m + \frac{1}{2} e^2 (1 - \cos 2m) + \frac{1}{8} e^3 (3 \cos m - 3 \cos 3m) \\ &\quad + \frac{1}{8} e^4 (\cos 4m - \cos 2m) + \text{etc.} \\ \therefore \cos v &= \cos m + \frac{1}{2} e (\cos 2m - 1) + (3e^2/8) (\cos 3m - \cos m) \\ &\quad + \frac{1}{8} e^3 (\cos 2m - \cos 4m) + \text{etc.} \end{aligned}$$

$$\begin{aligned} \therefore \sin v &= A \sin m + B \sin 2m + C \sin 3m + D \sin 4m + \dots \\ \cos v &= -\frac{1}{2} e + A_1 \cos m + B_1 \cos 2m + C_1 \cos 3m + D_1 \cos 4m + \dots \end{aligned}$$

where $A, B, C, D, \dots, A_1, B_1, C_1, D_1, \dots$ are each a series in powers of e .

$$\begin{aligned} \therefore x &= a \cos v = -\frac{1}{2} a e + a A_1 \cos m + a B_1 \cos 2m + a C_1 \cos 3m + a D_1 \cos 4m + \dots \\ y &= b \sin v = b A \sin m + b B \sin 2m + b C \sin 3m + b D \sin 4m + \dots \\ r &= a(1 - e \cos v) = a \left[1 + \frac{1}{2} e^2 - e A_1 \cos m - e B_1 \cos 2m - e C_1 \cos 3m \right. \\ &\quad \left. - e D_1 \cos 4m + \dots \right]. \end{aligned}$$

126. Proposed by J. SCHEFFER, A. M., Hagerstown, Md.

The declination of a certain fixed star is $12^\circ 40'$. Its altitude was observed one day to be $16^\circ 40'$. Three hours and twenty-four minutes later it was found to be $40^\circ 20'$. Find the latitude of the place of observation.

Solution by G. B. M. ZERR, A. M., Ph.D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Let $\delta=12^\circ 40'$, φ =latitude of observation, $a=16^\circ 40'$, $a'=40^\circ 20'$, $\mu=51^\circ=3$ hours, 24 minutes, h =hour angle.

$$\text{Then } \cosh = \frac{\sin a - \sin \varphi \sin \delta}{\cos \varphi \cos \delta} = x, \quad \cos(h - \mu) = \frac{\sin a' - \sin \varphi \sin \delta}{\cos \varphi \cos \delta} = y.$$

Eliminating h , $\sin^2 \mu = x^2 - 2xy \cos \mu + y^2$. Substituting values of x and y ,

$$\cos^2 \varphi \cos^2 \delta \sin^2 \mu = 2 \sin^2 \varphi \sin^2 \delta (1 - \cos \mu) + \sin^2 a + \sin^2 a' - 2 \sin a \sin a' \cos \mu - 2 \sin \varphi \sin \delta (1 - \cos \mu) (\sin a + \sin a').$$

$$\therefore \sin^2 \varphi [\sin^2 \mu + \sin^2 \delta (1 - \cos \mu)^2] - 2 \sin \varphi \sin \delta (1 - \cos \mu) (\sin a + \sin a') = \cos^2 \delta \sin^2 \mu - \sin^2 a - \sin^2 a' + 2 \sin a \sin a' \cos \mu.$$

$$\text{Let } \sin^2 \mu + \sin^2 \delta (1 - \cos \mu)^2 = A = .610569.$$

$$\sin \delta (1 - \cos \mu) (\sin a + \sin a') = B = .075921.$$

$$\cos^2 \delta \sin^2 \mu - \sin^2 a - \sin^2 a' + 2 \sin a \sin a' \cos \mu = C = .307394.$$

$$\therefore A \sin^2 \varphi - 2B \sin \varphi = C.$$

$$\therefore \sin \varphi = \frac{B \pm \sqrt{(AC + B^2)}}{A} = .844702, \text{ or } -.596013. \quad \varphi = 57^\circ 38' 37''.$$

PROBLEMS FOR SOLUTION.

ARITHMETIC.

166. Proposed by F. P. MATZ, Sc. D., Ph. D., Professor of Mathematics and Astronomy in Defiance College, Defiance, Ohio.

If I sell one of my farms for \$A,=\$4500, and the other for \$B,=\$1800, I will gain $p\%$, =5%, on cost of both; but if I sell the dearer farm for \$C,=\$4000, and the other at cost, I will lose $p\%$, =5%. Find the cost of each farm.

AVERAGE AND PROBABILITY.

138. Proposed by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics, The Temple College, Philadelphia, Pa.

Find the average area of (1) triangle, (2) quadrilateral, (3) pentagon, (4) hexagon, formed by taking (1) three, (2) four, (3) five, (4) six random points on the circumference of a given circle radius a .

139. Proposed by L. C. WALKER, A. M., Graduate Student, Leland Stanford Jr. University, Cal.

Four points are taken at random on the surface of a given sphere; find the average volume of the tetrahedron formed by the planes passing through the points taken three and three.